

# 演習問題 No. 6 の解答

[1] (1)

$$\int_{1/\sqrt{2}}^1 \frac{1}{\sqrt{2-x^2}} dx = \left[ \sin^{-1} \frac{x}{\sqrt{2}} \right]_{1/\sqrt{2}}^1 = \sin^{-1} \frac{1}{\sqrt{2}} - \sin^{-1} \frac{1}{2} = \frac{\pi}{4} - \frac{\pi}{6} = \frac{\pi}{12}$$

(2) 部分積分により

$$\begin{aligned} \int_1^4 \log(x+2) dx &= \int_1^4 (x+2)' \log(x+2) dx = \left[ (x+2) \log(x+2) \right]_1^4 - \int_1^4 (x+2) \frac{1}{x+2} dx \\ &= 6 \log 6 - 3 \log 3 - 3 = 6 \log 2 + 3 \log 3 - 3 \end{aligned}$$

(3)  $\tan^{-1} x = t$  とおけば,  $\frac{dt}{dx} = \frac{1}{1+x^2}$  より,

$$\int_0^{\sqrt{3}} \frac{\tan^{-1} x}{x^2+1} dx = \int_0^{\pi/3} t dt = \left[ \frac{t^2}{2} \right]_0^{\pi/3} = \frac{\pi^2}{18}$$

[2] (1)  $\sum_{k=1}^n \frac{1}{n+3k} = \sum_{k=1}^n \frac{1}{n} \frac{1}{1+3(k/n)}$  より, 関数  $\frac{1}{1+3x}$  を考えると,

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{1}{n+3k} = \int_0^1 \frac{1}{1+3x} dx = \left[ \frac{1}{3} \log(1+3x) \right]_0^1 = \frac{2}{3} \log 2$$

(2)  $\sum_{k=1}^n \frac{1}{\sqrt{3n^2+nk}} = \sum_{k=1}^n \frac{1}{n} \frac{1}{\sqrt{3+(k/n)}}$  より, 関数  $\frac{1}{\sqrt{3+x}}$  を考えると,

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{1}{\sqrt{n^2+nk}} = \int_0^1 \frac{1}{\sqrt{3+x}} dx = \left[ 2\sqrt{3+x} \right]_0^1 = 2(2-\sqrt{3})$$

[3] (1)  $x^2 = t$  とおけば,  $\frac{dt}{dx} = 2x$  より,

$$\int \frac{x}{x^4+1} dx = \int \frac{1}{t^2+1} \cdot \frac{1}{2} dt = \frac{1}{2} \tan^{-1} t = \frac{1}{2} \tan^{-1} x^2$$

(2)  $\sqrt{x} = t$  とおけば,  $x = t^2$ ,  $\frac{dx}{dt} = 2t$  より,

$$\int e^{\sqrt{x}} dx = 2 \int te^t dt = 2(t-1)e^t = 2(\sqrt{x}-1)e^{\sqrt{x}}$$

(3)

$$\int \frac{1}{\sqrt{6x-x^2}} dx = \int \frac{1}{\sqrt{9-(x-3)^2}} dx = \sin^{-1} \left( \frac{x-3}{3} \right)$$

(4) 部分積分により,

$$\begin{aligned} \int (\sin^{-1} x)^2 dx &= x(\sin^{-1} x)^2 - \int x 2 \sin^{-1} x \frac{1}{\sqrt{1-x^2}} dx \\ &= x(\sin^{-1} x)^2 - 2 \left( (-\sqrt{1-x^2}) \sin^{-1} x - \int (-\sqrt{1-x^2}) \frac{1}{\sqrt{1-x^2}} dx \right) \\ &= x(\sin^{-1} x)^2 + 2\sqrt{1-x^2} \sin^{-1} x - 2x \end{aligned}$$

(5)

$$\int \frac{1}{x^{10} + x} dx = \int \frac{1}{(1 + x^{-9})x^{10}} dx$$

より,  $1 + x^{-9} = t$  とおけば,  $\frac{dt}{dx} = -\frac{9}{x^{10}}$  となるから,

$$= -\frac{1}{9} \int \frac{1}{t} dt = -\frac{1}{9} \log |t| = -\frac{1}{9} \log \left| 1 + \frac{1}{x^9} \right|$$

4 (1)  $\int_0^\pi = \int_0^{\pi/2} + \int_{\pi/2}^\pi$  と分割し, 最後の積分に変数変換  $x = \pi - t$  を用いて,

$$\begin{aligned} \int_0^\pi xf(\sin x) dx &= \int_0^{\pi/2} xf(\sin x) dx + \int_{\pi/2}^0 (\pi - t)f(\sin(\pi - t))(-1) dt \\ &= \int_0^{\pi/2} xf(\sin x) dx + \pi \int_0^{\pi/2} f(\sin t) dt - \int_0^{\pi/2} tf(\sin t) dt \\ &= \pi \int_0^{\pi/2} f(\sin x) dx \end{aligned}$$

(2)  $\cos^2 x = 1 - \sin^2 x$  より, (1) の結果を用いて,

$$\int_0^\pi \frac{x \sin x}{1 + \cos^2 x} dx = \pi \int_0^{\pi/2} \frac{\sin x}{1 + \cos^2 x} dx = \pi \left[ -\tan^{-1}(\cos x) \right]_0^{\pi/2} = \pi \tan^{-1} 1 = \frac{\pi^2}{4}$$

5 変数変換  $9 - x = t$  を用いて,

$$I = \int_0^9 \frac{\sqrt{x}}{\sqrt{x} + \sqrt{9-x}} dx = \int_9^0 \frac{\sqrt{9-t}}{\sqrt{9-t} + \sqrt{t}} (-1) dt = \int_0^9 \frac{\sqrt{9-t}}{\sqrt{t} + \sqrt{9-t}} dt$$

となるから,

$$\begin{aligned} 2I &= \int_0^9 \frac{\sqrt{x}}{\sqrt{x} + \sqrt{9-x}} dx + \int_0^9 \frac{\sqrt{9-x}}{\sqrt{x} + \sqrt{9-x}} dx = \int_0^9 dx = 9 \\ I &= \int_0^9 \frac{\sqrt{x}}{\sqrt{x} + \sqrt{9-x}} dx = \frac{9}{2} \end{aligned}$$