


Meson-baryon interactions and baryon resonances



Tetsuo Hyodo

Tokyo Institute of Technology

Contents



$\Lambda(1405)$ in meson-baryon scattering

[T. Hyodo, D. Jido, arXiv:1104.4474, submitted to Prog. Part. Nucl. Phys.](#)

- Chiral SU(3) dynamics
- Pole structure of $\Lambda(1405)$



Toward realistic meson-baryon interaction

- Constraint by accurate $\bar{K}N$ data

[Y. Ikeda, T. Hyodo, W. Weise, in preparation](#)

- Information of $\pi\Sigma$ channel

[Y. Ikeda, T. Hyodo, D. Jido, H. Kamano, T. Sato, K. Yazaki, arXiv:1101.5190 \[nucl-th\], to appear in Prog. Theor. Phys., T. Hyodo, M. Oka, arXiv:1105.5494 \[nucl-th\]](#)



Summary

Chiral symmetry breaking in hadron physics

Chiral symmetry: QCD with massless quarks

Consequence of chiral symmetry breaking in hadron physics

- appearance of the Nambu-Goldstone (NG) boson

$$m_\pi \sim 140 \text{ MeV}$$

- dynamical generation of hadron masses

$$M_p \sim 1 \text{ GeV} \sim 3M_q, \quad M_q \sim 300 \text{ MeV} \quad v.s. \quad 3 - 7 \text{ MeV}$$

- constraints on the NG-boson--hadron interaction

low energy theorems \leftarrow current algebra

systematic low energy ($m, p/4\pi f_\pi$) expansion: ChPT

Chiral symmetry and its breaking


$$SU(3)_R \otimes SU(3)_L \rightarrow SU(3)_V$$

Underlying QCD \Leftrightarrow observed hadron phenomena

s-wave low energy interaction in ChPT

Leading order term for the meson-baryon scattering

$$\mathcal{L}^{\text{WT}} = \frac{1}{4f^2} \text{Tr} \left(\bar{B} i \gamma^\mu [\Phi \partial_\mu \Phi - (\partial_\mu \Phi) \Phi], B \right)$$



s-wave contribution: Weinberg-Tomozawa (WT) term

Y. Tomozawa, *Nuovo Cim.* **46A**, 707 (1966); S. Weinberg, *Phys. Rev. Lett.* **17**, 616 (1966)

$$V_{ij} = -\frac{C_{ij}}{4f^2} (\omega_i + \omega_j) + \dots$$

$$C_{ij} = \sum_{\alpha} [6 - C_2(\alpha)] \left(\begin{array}{cc} 8 & 8 \\ I_{\bar{i}}, Y_{\bar{i}} & I_i, Y_i \end{array} \parallel \begin{array}{c} \alpha \\ I, Y \end{array} \right) \left(\begin{array}{cc} 8 & 8 \\ I_{\bar{j}}, Y_{\bar{j}} & I_j, Y_j \end{array} \parallel \begin{array}{c} \alpha \\ I, Y \end{array} \right)$$

$$Y = Y_{\bar{i}} + Y_i = Y_{\bar{j}} + Y_j, \quad I = I_{\bar{i}} + I_i = I_{\bar{j}} + I_j,$$

- Flavor SU(3) symmetry --> **sign and strength**
- Derivative coupling --> **energy dependence**
- Systematic improvement by higher order terms (later)

If the interaction is strong, resummation is mandatory.

Scattering amplitude and unitarity

Unitarity of S-matrix: Optical theorem

$$\text{Im}[T^{-1}(s)] = \frac{\rho(s)}{2} \leftarrow \text{phase space of two-body state}$$

General amplitude by dispersion relation

$$T^{-1}(\sqrt{s}) = \sum_i \frac{R_i}{\sqrt{s} - W_i} + \tilde{a}(s_0) + \frac{s - s_0}{2\pi} \int_{s_+}^{\infty} ds' \frac{\rho(s')}{(s' - s)(s' - s_0)}$$

R_i, W_i, a : to be determined by chiral interaction

Identify dispersion integral = loop function G , the rest = V^{-1}

$$T(\sqrt{s}) = \frac{1}{V^{-1}(\sqrt{s}) - G(\sqrt{s}; a)}$$

Scattering amplitude

The function V is determined by the **matching with ChPT**

$$T^{(1)} = V^{(1)}, \quad T^{(2)} = V^{(2)}, \quad T^{(3)} = V^{(3)} - V^{(1)}GV^{(1)}, \quad \dots$$

Amplitude T : consistent with chiral symmetry + unitarity

Chiral unitary approach


Meson-baryon scattering amplitude

- Interaction \leftarrow chiral symmetry

Y. Tomozawa, *Nuovo Cim.* 46A, 707 (1966); S. Weinberg, *Phys. Rev. Lett.* 17, 616 (1966)

- Amplitude \leftarrow unitarity in **coupled channels**

R.H. Dalitz, T.C. Wong, G. Rajasekaran, *Phys. Rev.* 153, 1617 (1967)



N. Kaiser, P. B. Siegel, W. Weise, *Nucl. Phys.* A594, 325 (1995),

E. Oset, A. Ramos, *Nucl. Phys.* A635, 99 (1998),

J. A. Oller, U. G. Meissner, *Phys. Lett.* B500, 263 (2001),

M.F.M. Lutz, E. E. Kolomeitsev, *Nucl. Phys.* A700, 193 (2002), ... many others

It successfully reproduces the scattering observables as well as the dynamically generated resonances.

The $\Lambda(1405)$ resonance


$\Lambda(1405) : J^P = 1/2^-, I = 0$

(PDG)

mass : 1406.5 ± 4.0 MeV, width : 50 ± 2 MeV

decay mode: $\Lambda(1405) \rightarrow (\pi\Sigma)_{I=0}$ **100%**

“naive” quark model
: p-wave ~ 1600 MeV?



N. Isgur, G. Karl, PRD18, 4187 (1978)

M B Coupled channel
multi-scattering




R.H. Dalitz, T.C. Wong,
G. Rajasekaran, PR153, 1617 (1967)

$\bar{K}N$ interaction below threshold

T. Hyodo, W. Weise, PRC 77, 035204 (2008)







--> $\bar{K}N$ potential, kaonic nuclei

A. Dote, T. Hyodo, W. Weise,
NPA804, 197 (2008); PRC 79, 014003 (2009)



A simple model (1 parameter) v.s. experimental data


Total cross section of K-p scattering



Branching ratio

	γ	R_c	R_n
exp.	2.36	0.664	0.189
theo.	1.80	0.624	0.225

$\pi\Sigma$ spectrum



T. Hyodo, S.I. Nam, D. Jido, A. Hosaka, PRC68, 018201 (2003); PTP 112, 73 (2004)

Good agreement with data above, at, and below $\bar{K}N$ threshold
more quantitatively --> fine tuning, higher order terms,...

Pole structure in the complex energy plane

Resonance state \sim pole of the scattering amplitude

D. Jido, J.A. Oller, E. Oset, A. Ramos, U.G. Meissner, Nucl. Phys. A 723, 205 (2003)


$$T_{ij}(\sqrt{s}) \sim \frac{g_i g_j}{\sqrt{s} - M_R + i\Gamma_R/2}$$

\sim

Two poles for one resonance (bump structure)

--> Superposition of two states ?

--> different $\pi\Sigma$ spectra?



T. Hyodo, D. Jido, arXiv:1104.4474

What is the **origin** of this structure?

Origin of the two-pole structure

Leading order chiral interaction for $\bar{K}N$ - $\pi\Sigma$ channel

T. Hyodo, W. Weise, Phys. Rev. C 77, 035204 (2008)


$$V_{ij} = -C_{ij} \frac{\omega_i + \omega_j}{4f^2}$$

$$C_{ij} = \begin{pmatrix} \bar{K}N & \pi\Sigma \\ 3 & -\sqrt{\frac{3}{2}} \\ -\sqrt{\frac{3}{2}} & 4 \end{pmatrix}$$

at threshold

$$\omega_i \sim m_i, \quad 3.3m_\pi \sim m_K$$

$$\Rightarrow V_{\bar{K}N} \sim 2.5V_{\pi\Sigma}$$



Very strong attraction in $\bar{K}N$ (higher energy) --> bound state
Strong attraction in $\pi\Sigma$ (lower energy) --> resonance


Model dependence? Effects from higher order terms?

Experimental constraints for $S=-1$ MB scattering

K-p total cross sections (bubble chamber, large errors)

$\bar{K}N$ threshold observables

- threshold branching ratios (old but accurate)
- K-p scattering length \leftarrow **SIDDHARTA exp.**



$\pi\Sigma$ mass spectra

- new data is becoming available (LEPS, CLAS, HADES,...)
- normalization, reaction dependence,... \leftarrow to be predicted?

$\pi\Sigma$ threshold observables (so far no data)

Constraints from $\bar{K}N$ data

- K-p total cross sections

$$K^- p \rightarrow (K^- p, \bar{K}^0 n, \pi^0 \Lambda, \pi^0 \Sigma^0, \pi^+ \Sigma^-, \pi^- \Sigma^+)$$

- Threshold branching ratios

$$\gamma = \frac{\Gamma(K^- p \rightarrow \pi^+ \Sigma^-)}{\Gamma(K^- p \rightarrow \pi^- \Sigma^+)} = 2.36 \pm 0.04$$

$$R_c = \frac{\Gamma(K^- p \rightarrow \pi^+ \Sigma^-, \pi^- \Sigma^+)}{\Gamma(K^- p \rightarrow \text{all inelastic channels})} = 0.664 \pm 0.011$$

$$R_n = \frac{\Gamma(K^- p \rightarrow \pi^0 \Lambda)}{\Gamma(K^- p \rightarrow \text{neutral states})} = 0.189 \pm 0.015$$

R.J. Nowak, *et al.*, Nucl. Phys. B139, 61 (1978); D.N. Tovee, *et al.*, *ibid*, B33, 493 (1971)


- Shift and width of 1s level of kaonic hydrogen (SIDDHARTA)

$$\Delta E = -283 \pm 36 \pm 6 \text{ eV}, \quad \Gamma = 541 \pm 89 \pm 22 \text{ eV}$$

Bazzi, *et al.*, arXiv:1105.3090 [nucl-ex]

$$\Delta E - \frac{i}{2}\Gamma = -2\alpha^3 \mu_c^2 a_{K^- p} [1 - 2\alpha \mu_c (\ln \alpha - 1) a_{K^- p}] \leftarrow \text{scattering length}$$

U.-G. Meissner, U. Raha, A. Rusetsky, Eur. Phys. J. C35, 349 (2004)



Construction of the realistic amplitude

Systematic χ^2 fitting with SIDDHARTA data


Y. Ikeda, T. Hyodo, W. Weise, in preparation

Interaction kernel: NLO ChPT

B. Borasoy, R. Nissler, W. Weise, *Eur. Phys. J. A25*, 79-96 (2005);

B. Borasoy, U.G. Meissner, R. Nissler, *Phys. Rev. C74*, 055201 (2006)


1) WT term



$\mathcal{O}(p)$

WT model


2) Born terms



$\mathcal{O}(p)$

WTB model

3) NLO terms




$\mathcal{O}(p^2)$

NLO model

Parameters: 6 cutoffs (+ 7 low energy constants in NLO)

Result by NLO model

Observables	Theory	Experiment
ΔE (eV)	306	283 ± 42
Γ (eV)	591	541 ± 111
γ	2.36	2.36 ± 0.04
R_c	0.659	0.664 ± 0.011
R_n	0.192	0.189 ± 0.015
a_{K-p} (fm)	$-0.81 + i0.87$	
Poles of the $\Lambda(1405)$ (MeV)	$1424.2 - i26.3, 1380.7 - i81.3$	




Good description of data ($\chi^2/\text{dof} \sim 1$)

Summary of results


Results from three models

	WT	WTB	NLO
χ^2/dof	1.12	1.15	0.957

Shift and width



Pole positions



Error analysis is now underway

Y. Ikeda, T. Hyodo, W. Weise, in preparation

$\pi\Sigma$ threshold behavior

Effect of the $\pi\Sigma$ threshold data for $\bar{K}N$ - $\pi\Sigma$ amplitude



Y. Ikeda, T. Hyodo, D. Jido, H. Kamano, T. Sato, K. Yazaki,
arXiv:1101.5190 [nucl-th], to appear in Prog. Theor. Phys.

Extrapolations with a given $\bar{K}N(l=0)$ scattering length
 --> uncertainty in subthreshold

Model	A1	A2	B E-dep	B E-indep
parameter ($\pi\Sigma$)	$d_{\pi\Sigma} = -1.67$	$d_{\pi\Sigma} = -2.85$	$\Lambda_{\pi\Sigma} = 1005$ MeV	$\Lambda_{\pi\Sigma} = 1465$ MeV
parameter ($\bar{K}N$)	$d_{\bar{K}N} = -1.79$	$d_{\bar{K}N} = -2.05$	$\Lambda_{\bar{K}N} = 1188$ MeV	$\Lambda_{\bar{K}N} = 1086$ MeV
pole 1 [MeV]	$1422 - 16i$	$1425 - 11i$	$1422 - 22i$	$1423 - 29i$
pole 2 [MeV]	$1375 - 72i$ (R)	1321 (B)	$1349 - 54i$ (R)	1325 (V)
$a_{\pi\Sigma}$ [fm]	0.934	-2.30	1.44	5.50
r_e [fm]	5.02	5.89	3.96	0.458
$a_{\bar{K}N}$ [fm] (input)	$-1.70 + 0.68i$	$-1.70 + 0.68i$	$-1.70 + 0.68i$	$-1.70 + 0.68i$

subthreshold behavior

← $\pi\Sigma$ scattering length,
 effective range



Determination of the $\pi\Sigma$ scattering length

$\pi\pi$ scattering length from $K \rightarrow \pi\pi\pi$ decay



N. Cabibbo, *Phys. Rev. Lett.* **93**, 121801 (2004);

NA48/2, J.R. Batley, *et al.*, *Phys. Lett. B* **686**, 101 (2010)

Analogy: $\pi\Sigma$ scattering lengths from $\Lambda_c \rightarrow \pi\pi\Sigma$ decays

T. Hyodo, M. Oka, arXiv:1105.5494 [nucl-th]


- isospin violation
- + threshold cusp
- + amplitude interference



Expansion of the spectrum around cusp \rightarrow scattering length

Summary 1

We study the $\bar{K}N$ - $\pi\Sigma$ interaction and $\Lambda(1405)$ based on chiral SU(3) symmetry and unitarity

 Chiral symmetry constrains the **NG boson dynamics** with hadrons.


 Unitarity should be taken into account for a **strongly interacting** system.

 **Two poles** for $\Lambda(1405)$


<-- attractive **$\bar{K}N$ and $\pi\Sigma$ interactions**

Summary 2

Recent developments to construct a realistic meson-baryon interaction

-  New $\bar{K}N$ threshold data by SIDDAHRTA
 - systematic χ^2 analysis with NLO terms

Y. Ikeda, T. Hyodo, W. Weise, in preparation

-  Threshold information of $\pi\Sigma$ channel
 - importance of $\pi\Sigma$ threshold behavior

Y. Ikeda, T. Hyodo, D. Jido, H. Kamano, T. Sato, K. Yazaki,
arXiv:1101.5190 [nucl-th], to appear in Prog. Theor. Phys.

- scattering length from Λ_c decay

T. Hyodo, M. Oka, arXiv:1105.5494 [nucl-th]