

Compositeness of hadrons from effective field theory



Tetsuo Hyodo

Yukawa Institute for Theoretical Physics, Kyoto Univ.

2016, Aug. 17th 1

Contents




Introduction

- Hadron structure, exotics, and clustering
- Resonance in hadron physics



Compositeness of near-threshold resonances

- Weak binding relation from EFT
- Generalization to resonances
- Application: $\Lambda(1405)$



T. Hyodo, Int. J. Mod. Phys. A 28, 1330045 (2013)

Y. Kamiya, T. Hyodo, Phys. Rev. C93, 035203 (2016), arXiv:1607.01899[hep-ph]


Y. Tsuchida, T. Hyodo, in preparation

**Talks by Y. Kamiya,
Y. Tsuchida on 18th**

Classification of hadrons

Observed hadrons

p	1/2 ⁺ ****	$\Delta(1232)$	3/2 ⁺ ****	Σ^+	1/2 ⁺ ****	Ξ^0	1/2 ⁺ ****	Λ_c^+	1/2 ⁺ ****
n	1/2 ⁺ ****	$\Delta(1600)$	3/2 ⁺ ***	Σ^0	1/2 ⁺ ****	Ξ^-	1/2 ⁺ ****	$\Lambda_c(2595)^+$	1/2 ⁺ ***
$N(1440)$	1/2 ⁺ ****	$\Delta(1620)$	1/2 ⁻ ****	Σ^-	1/2 ⁺ ****	$\Xi(1530)$	3/2 ⁺ ****	$\Lambda_c(2625)^+$	3/2 ⁻ ***
$N(1520)$	3/2 ⁻ ***	$\Delta(1700)$	3/2 ⁻ ***	$\Sigma(1385)$	3/2 ⁺ ***	$\Xi(1620)$	*	$\Lambda_c(2765)^+$	*
$N(1535)$	1/2 ⁻ ***	$\Delta(1750)$	1/2 ⁺ *	$\Sigma(1480)$	*	$\Xi(1690)$	***	$\Lambda_c(2880)^+$	5/2 ⁺ ***
$N(1650)$	1/2 ⁻ ***	$\Delta(1900)$	1/2 ⁻ **	$\Sigma(1560)$	**	$\Xi(1820)$	3/2 ⁻ ***	$\Lambda_c(2940)^+$	***
$N(1675)$	5/2 ⁻ ***	$\Delta(1905)$	5/2 ⁺ ****	$\Sigma(1580)$	3/2 ⁻ *	$\Xi(1950)$	***	$\Sigma_c(2455)$	1/2 ⁺ ****
$N(1680)$	5/2 ⁺ ***	$\Delta(1910)$	1/2 ⁺ ****	$\Sigma(1620)$	1/2 ⁻ *	$\Xi(2030)$	$\geq \frac{5}{2}$ ***	$\Sigma_c(2520)$	3/2 ⁺ ***
$N(1685)$	*	$\Delta(1920)$	3/2 ⁺ ***	$\Sigma(1660)$	1/2 ⁺ ***	$\Xi(2120)$	*	$\Sigma_c(2800)$	***
$N(1700)$	3/2 ⁻ ***	$\Delta(1930)$	5/2 ⁻ ***	$\Sigma(1670)$	3/2 ⁻ ***	$\Xi(2250)$	**	$\Xi_c(2645)$	3/2 ⁺ ***
$N(1710)$	1/2 ⁺ ***	$\Delta(1940)$	3/2 ⁻ **	$\Sigma(1690)$	**	$\Xi(2370)$	**	$\Xi_c(2790)$	1/2 ⁻ ***
$N(1720)$	3/2 ⁺ ***	$\Delta(1950)$	7/2 ⁺ ****	$\Sigma(1730)$	3/2 ⁺ *	$\Xi(2500)$	*	$\Xi_c(2815)$	3/2 ⁻ ***
$N(1860)$	5/2 ⁺ **	$\Delta(2000)$	5/2 ⁺ **	$\Sigma(1750)$	1/2 ⁻ ***	$\Xi(2645)$	3/2 ⁺ ***	$\Xi_c(2930)$	*
$N(1875)$	3/2 ⁻ ***	$\Delta(2150)$	1/2 ⁻ *	$\Sigma(1770)$	1/2 ⁺ *	Ω^-	3/2 ⁺ ****	$\Xi_c(2980)$	***
$N(1880)$	1/2 ⁺ **	$\Delta(2200)$	7/2 ⁻ *	$\Sigma(1775)$	5/2 ⁻ ***	$\Omega(2250)^-$	***	$\Xi_c(3055)$	***
$N(1895)$	1/2 ⁻ **	$\Delta(2300)$	9/2 ⁺ **	$\Sigma(1840)$	3/2 ⁺ *	$\Omega(2380)^-$	**	$\Xi_c(3080)$	***
$N(1900)$	3/2 ⁺ ***	$\Delta(2350)$	5/2 ⁻ *	$\Sigma(1880)$	1/2 ⁺ **	$\Omega(2470)^-$	**	$\Xi_c(3123)$	*
$N(1990)$	7/2 ⁺ **	$\Delta(2390)$	7/2 ⁺ *	$\Sigma(1900)$	1/2 ⁻ *	Ω_c^0	1/2 ⁺ ***	$\Xi_c(2980)$	***
$N(2000)$	5/2 ⁺ **	$\Delta(2400)$	9/2 ⁻ **	$\Sigma(1915)$	5/2 ⁺ ****	$\Omega_c(2770)^0$	3/2 ⁺ ***	$\Xi_c(3055)$	***
$N(2040)$	3/2 ⁺ *	$\Delta(2420)$	11/2 ⁺ ****	$\Sigma(1940)$	3/2 ⁺ *	$\Xi_c(2770)^0$	3/2 ⁺ ***	$\Xi_c(3123)$	*
$N(2060)$	5/2 ⁻ **	$\Delta(2750)$	13/2 ⁻ **	$\Sigma(1940)$	3/2 ⁻ ***	$\Xi_c(2770)^0$	3/2 ⁺ ***	$\Xi_c(3123)$	*
$N(2100)$	1/2 ⁺ *	$\Delta(2950)$	15/2 ⁺ **	$\Sigma(2000)$	1/2 ⁻ *	$\Xi_c(2770)^0$	3/2 ⁺ ***	$\Xi_c(2770)^0$	3/2 ⁺ ***
$N(2120)$	3/2 ⁻ **			$\Sigma(2030)$	7/2 ⁺ ****	$\Xi_c(2770)^0$	3/2 ⁺ ***	$\Xi_c(2770)^0$	3/2 ⁺ ***
$N(2190)$	7/2 ⁻ ***	Λ	1/2 ⁺ ****	$\Sigma(2070)$	5/2 ⁺ *	Ξ_{cc}^+	*	$\Xi_c(2770)^0$	3/2 ⁺ ***
$N(2220)$	9/2 ⁻ ***	$\Lambda(1405)$	1/2 ⁻ ***	$\Sigma(2080)$	3/2 ⁺ **			$\Xi_c(2770)^0$	3/2 ⁺ ***
$N(2250)$	9/2 ⁻ ***	$\Lambda(1520)$	3/2 ⁻ ***	$\Sigma(2100)$	7/2 ⁻ *			$\Xi_c(2770)^0$	3/2 ⁺ ***
$N(2300)$	1/2 ⁺ **	$\Lambda(1600)$	1/2 ⁺ ***	$\Sigma(2250)$	***			$\Xi_c(2770)^0$	3/2 ⁺ ***
$N(2570)$	5/2 ⁻ **	$\Lambda(1670)$	1/2 ⁻ ***	$\Sigma(2455)$	**			$\Xi_c(2770)^0$	3/2 ⁺ ***
$N(2600)$	11/2 ⁻ ***	$\Lambda(1690)$	3/2 ⁻ ***	$\Sigma(2620)$	**			$\Xi_c(2770)^0$	3/2 ⁺ ***
$N(2700)$	13/2 ⁺ **	$\Lambda(1710)$	1/2 ⁺ *	$\Sigma(3000)$	*			$\Xi_c(2770)^0$	3/2 ⁺ ***
$\Lambda(1800)$	1/2 ⁻ ***			$\Sigma(3170)$	*			$\Xi_c(2770)^0$	3/2 ⁺ ***
$\Lambda(1810)$	1/2 ⁺ ***							$\Xi_c(2770)^0$	3/2 ⁺ ***
$\Lambda(1820)$	5/2 ⁻ ***							$\Xi_c(2770)^0$	3/2 ⁺ ***
$\Lambda(1830)$	5/2 ⁻ ***							$\Xi_c(2770)^0$	3/2 ⁺ ***
$\Lambda(1890)$	3/2 ⁺ ***							$\Xi_c(2770)^0$	3/2 ⁺ ***
$\Lambda(2000)$	*							$\Xi_c(2770)^0$	3/2 ⁺ ***
$\Lambda(2020)$	7/2 ⁺ *							$\Xi_c(2770)^0$	3/2 ⁺ ***
$\Lambda(2050)$	3/2 ⁻ *							$\Xi_c(2770)^0$	3/2 ⁺ ***
$\Lambda(2100)$	7/2 ⁻ ***							$\Xi_c(2770)^0$	3/2 ⁺ ***
$\Lambda(2110)$	5/2 ⁺ ***							$\Xi_c(2770)^0$	3/2 ⁺ ***
$\Lambda(2325)$	3/2 ⁻ *							$\Xi_c(2770)^0$	3/2 ⁺ ***
$\Lambda(2350)$	9/2 ⁺ ***							$\Xi_c(2770)^0$	3/2 ⁺ ***
$\Lambda(2585)$	**							$\Xi_c(2770)^0$	3/2 ⁺ ***



~ 150 baryons

PDG2015 : <http://pdg.lbl.gov/>

LIGHT UNFLAVORED ($S = C = B = 0$)		STRANGE ($S = \pm 1, C = B = 0$)		CHARMED, STRANGE ($C = S = \pm 1$)		$\Xi_c(F^C)$	
$F(F^C)$	$\bar{F}(F^C)$	$K(F^C)$	$\bar{K}(F^C)$	$K(F^C)$	$\bar{K}(F^C)$	$K(F^C)$	$\bar{K}(F^C)$
$\bullet \pi^\pm$	1 ⁻ (0 ⁻)	$\bullet \phi(1680)$	0 ^{+(1⁻)}	$\bullet K^\pm$	1/2(0 ⁻)	$\bullet D_s^\pm$	0(0 ⁻)
$\bullet \pi^0$	1 ⁻ (0 ⁻)	$\bullet \rho(1690)$	1 ^{+(3⁻)}	$\bullet K^0$	1/2(0 ⁻)	$\bullet D_s^{\pm\pm}$	0(?)
$\bullet f_0(500)$	0 ^{+(0⁺)}	$\bullet \rho(1700)$	1 ^{+(1⁻)}	$\bullet K_S^0$	1/2(0 ⁻)	$\bullet D_S^0(2317)^\pm$	0(0 ⁺)
$\bullet \pi(770)$	1 ^{+(1⁻)}	$\bullet \rho(1710)$	1 ^{+(2⁺)}	$\bullet K_1^0$	1/2(0 ⁻)	$\bullet D_S(2460)^\pm$	0(1 ⁺)
$\bullet \omega(782)$	0 ^{-(1⁻)}	$\bullet \eta(1760)$	0 ^{+(0⁺)}	$\bullet K'(892)$	1/2(1 ⁻)	$\bullet D_S(2536)^\pm$	0(1 ⁺)
$\bullet \pi(958)$	0 ^{+(0⁺)}	$\bullet \pi(1800)$	1 ^{+(0⁻)}	$\bullet K(1270)$	1/2(2 ⁻)	$\bullet D_S(2700)^\pm$	0(1 ⁻)
$\bullet f_0(980)$	0 ^{+(0⁺)}	$\bullet f_0(1810)$	0 ^{+(2⁺)}	$\bullet K(1400)$	1/2(2 ⁺)	$\bullet D_s^*(2860)^\pm$	0(?)
$\bullet \omega(1020)$	0 ^{-(1⁻)}	$\bullet \chi(1840)$?	$\bullet K'(1410)$	1/2(2 ⁻)	$D_s(3040)^\pm$	0(?)
$\bullet h_1(1170)$	0 ^{-(1⁻)}	$\bullet \phi(1850)$	0 ^{-(3⁻)}	$\bullet K(1430)$	1/2(2 ⁺)	BOTTOM ($B = \pm 1$)	
$\bullet b_1(1235)$	1 ^{+(1⁻)}	$\bullet \pi(1870)$	0 ^{+(2⁺)}	$\bullet K(1460)$	1/2(0 ⁻)	ADMIXTURE	
$\bullet a_1(1260)$	1 ^{-(1⁺)}	$\bullet \pi(1880)$	1 ^{-(2⁻)}	$\bullet K(1580)$	1/2(2 ⁻)	$\bullet B_s^{\pm}$	1/2(0 ⁻)
$\bullet f_0(1270)$	0 ^{+(2⁺)}	$\bullet \rho(1900)$	1 ^{+(1⁻)}	$\bullet K(1650)$	1/2(1 ⁻)	$\bullet B_s^0$	1/2(0 ⁻)
$\bullet f_0(1285)$	0 ^{+(0⁺)}	$\bullet \pi(1910)$	0 ^{+(2⁺)}	$\bullet K(1660)$	1/2(1 ⁻)	$\bullet B_s^0/B_s^0/B_s^0$	Baryon
$\bullet (\pi)(1300)$	1 ^{-(0⁻)}	$\bullet \rho(1920)$	0 ^{+(2⁺)}	$\bullet K(1770)$	1/2(2 ⁻)	$\bullet K(1770)$	ADMIXTURE
$\bullet a_2(1320)$	1 ^{-(2⁺)}	$\bullet \pi(1930)$	0 ^{+(0⁺)}	$\bullet K(1780)$	1/2(3 ⁻)	$\bullet V_u$ and V_b	CKM Matrix Elements
$\bullet f_0(1370)$	0 ^{+(0⁺)}	$\bullet f_0(1940)$	0 ^{+(1⁻)}	$\bullet K(1820)$	1/2(2 ⁻)	$\bullet B_s^0$	1/2(1 ⁻)
$\bullet h_1(1380)$?	$\bullet a_1(1940)$	1 ^{-(4⁺)}	$\bullet K(1830)$	1/2(0 ⁻)	$\bullet B_s(5721)^\pm$	1/2(1 ⁻)
$\bullet \pi_1(1400)$	1 ^{-(1⁻)}	$\bullet \pi_1(1950)$	0 ^{+(4⁺)}	$\bullet K(1950)$	1/2(2 ⁺)	$\bullet B_s(5721)^\pm$?
$\bullet (\pi_1)(1405)$	0 ^{+(0⁺)}	$\bullet \pi_2(1960)$	1 ^{-(2⁻)}	$\bullet K(1980)$	1/2(2 ⁺)	$\bullet B_s(5721)^\pm$?
$\bullet f_1(1420)$	0 ^{+(1⁺)}	$\bullet f_1(1970)$	0 ^{+(1⁺)}	$\bullet K(2010)$	1/2(2 ⁺)	$\bullet B_s(5747)^\pm$	1/2(2 ⁺)
$\bullet f_2(1420)$	0 ^{-(1⁻)}	$\bullet f_2(1970)$	0 ^{+(2⁺)}	$\bullet K(2020)$	1/2(2 ⁻)	$\bullet B_s(5747)^\pm$	1/2(2 ⁺)
$\bullet f_0(1430)$	0 ^{+(2⁺)}	$\bullet \rho(1970)$	1 ^{-(1⁻)}	$\bullet K(2150)$	1/2(1 ⁻)	$\bullet B_s(5770)^\pm$?
$\bullet a_1(1450)$	1 ^{-(0⁺)}	$\bullet \pi(1980)$	1 ^{-(1⁻)}	$\bullet K(2200)$	1/2(2 ⁻)	$\bullet B_s(5770)^\pm$?
$\bullet (\pi_1)(1475)$	0 ^{+(0⁺)}	$\bullet \pi_2(1990)$	1 ^{-(1⁻)}	$\bullet K(2200)$	1/2(2 ⁻)	$\bullet B_s(5770)^\pm$?
$\bullet f_0(1500)$	0 ^{+(0⁺)}	$\bullet \eta(2020)$	1 ^{-(3⁻)}	$\bullet K(2300)$	1/2(3 ⁻)	$\bullet B_s^0$	0(0 ⁻)
$\bullet f_0(1525)$	0 ^{+(2⁺)}	$\bullet \rho(2030)$	0 ^{+(0⁺)}	$\bullet f_0(2300)$	1/2(4 ⁻)	$\bullet B_s^0$	0(1 ⁺)
$\bullet f_0(1565)$	0 ^{+(2⁺)}	$\bullet \rho(2040)$	0 ^{+(0⁺)}	$\bullet f_0(2300)$	1/2(4 ⁻)	$\bullet B_s^0(5830)^\pm$	0(1 ⁺)
$\bullet f_0(1570)$	1 ^{+(1⁻)}	$\bullet \rho(2040)$	0 ^{+(0⁺)}	$\bullet f_0(2300)$	1/2(4 ⁻)	$\bullet B_s^0(5840)^\pm$	0(2 ⁺)
$\bullet h_1(1595)$	0 ^{-(1⁺)}	$\bullet \rho(2040)$	0 ^{+(2⁺)}	$\bullet f_0(2340)$	1/2(4 ⁻)	$\bullet B_s^0(5850)^\pm$?
$\bullet \pi_1(1600)$	1 ^{-(1⁻)}	$\bullet \rho(2050)$	1 ^{-(6⁻)}	$\bullet \rho_1(2050)$	1/2(4 ⁻)	$\bullet D_s^0(2400)^\pm$	1/2(0 ⁻)
$\bullet \rho_1(2350)$	1 ^{-(1⁻)}	$\bullet \rho_2(2060)$	1 ^{-(6⁻)}	$\bullet \rho_2(2060)$	1/2(2 ⁻)	$\bullet D_s^0(2420)^\pm$	1/2(2 [?])
$\bullet \rho_2(2450)$	1 ^{-(6⁻)}	$\bullet \rho_2(2070)$	1 ^{-(6⁻)}	$\bullet \rho_2(2070)$	1/2(2 ⁻)	$\bullet D_s^0(2420)^\pm$	1/2(2 [?])
$\bullet \rho_2(2600)$	1 ^{-(6⁻)}	$\bullet \rho_2(2080)$	1 ^{-(6⁻)}	$\bullet \rho_2(2080)$	1/2(2 ⁻)	$\bullet D_s^0(2430)^\pm$	1/2(2 [?])
$\bullet \rho_2(2750)$	1 ^{-(6⁻)}	$\bullet \rho_2(2090)$	1 ^{-(6⁻)}	$\bullet \rho_2(2090)$	1/2(2 ⁻)	$\bullet D_s^0(2430)^\pm$	1/2(2 [?])
$\bullet \rho_2(2750)$	1 ^{-(6⁻)}	$\bullet \rho_2(2100)$	1 ^{-(6⁻)}	$\bullet \rho_2(2100)$	1/2(2 ⁻)	$\bullet D_s^0(2430)^\pm$	1/2(2 [?])
$\bullet \rho_2(2750)$	1 ^{-(6⁻)}	$\bullet \rho_2(2110)$	1 ^{-(6⁻)}	$\bullet \rho_2(2110)$	1/2(2 ⁻)	$\bullet D_s^0(2430)^\pm$	1/2(2 [?])
$\bullet \rho_2(2750)$	1 ^{-(6⁻)}	$\bullet \rho_2(2120)$	1 ^{-(6⁻)}	$\bullet \rho_2(2120)$	1/2(2 ⁻)	$\bullet D_s^0(2430)^\pm$	1/2(2 [?])
$\bullet \rho_2(2750)$	1 ^{-(6⁻)}	$\bullet \rho_2(2130)$	1 ^{-(6⁻)}	$\bullet \rho_2(2130)$	1/2(2 ⁻)	$\bullet D_s^0(2430)^\pm$	1/2(2 [?])
$\bullet \rho_2(2750)$	1 ^{-(6⁻)}	$\bullet \rho_2(2140)$	1 ^{-(6⁻)}	$\bullet \rho_2(2140)$	1/2(2 ⁻)	$\bullet D_s^0(2430)^\pm$	1/2(2 [?])
$\bullet \rho_2(2750)$	1 ^{-(6⁻)}	$\bullet \rho_2(2150)$	1 ^{-(6⁻)}	$\bullet \rho_2(2150)$	1/2(2 ⁻)	$\bullet D_s^$	


States with more than $qqq/q\bar{q}$

Tetraquark candidate (Belle)

: $Z_b(10610), Z_b(10650)$

$$\begin{aligned} Y(5S) \rightarrow & \pi^\pm + Z_b \\ \hookrightarrow & Y(nS)(b\bar{b}) + \pi^\mp(u\bar{d}/d\bar{u}) \end{aligned}$$

A. Bondar, *et al.*, Phys. Rev. Lett. 108, 122001 (2012)




Pentaquark candidate (LHCb)

: $P_c(4450), P_c(4380)$

$$\begin{aligned} \Lambda_b \rightarrow & K^- + P_c \\ \hookrightarrow & J/\psi(c\bar{c}) + p(uud) \end{aligned}$$

R. Aaij, *et al.*, Phys. Rev. Lett. 115, 072001 (2015)




Only a few are observed. Why only a few?


Various hadronic excitations

Description of excited baryons


Conventional structure



Exotic structures



multiquark
q \bar{q} pair creation



hadronic
molecule
(clustering)


In QCD, non-qqq structures naturally arise.

- Baryons: superposition of qqq + exotic structures
- > How can we disentangle different components?

Unstable states via strong interaction

Hadron resonances

p	1/2 ⁺ ****	$\Delta(1232)$	3/2 ⁺ ****	Σ^+	1/2 ⁺ ****	Ξ^0	1/2 ⁺ ****	Λ_c^+	1/2 ⁺ ****
n	1/2 ⁺ ****	$\Delta(1600)$	3/2 ⁺ ***	Σ^0	1/2 ⁺ ****	Ξ^-	1/2 ⁺ ****	$\Lambda_c(2595)^+$	1/2 ⁻ ***
$N(1440)$	1/2 ⁺ ****	$\Delta(1620)$	1/2 ⁻ ***	$\Sigma^-(1385)$	3/2 ⁺ ****	$\Xi(1530)$	3/2 ⁺ ****	$\Lambda_c(2625)^+$	3/2 ⁻ ***
$N(1520)$	3/2 ⁻ ***	$\Delta(1700)$	3/2 ⁻ ***	$\Sigma(1480)$	*	$\Xi(1620)$	*	$\Lambda_c(2765)^+$	*
$N(1535)$	1/2 ⁻ ***	$\Delta(1750)$	1/2 ⁺ *	$\Sigma(1560)$	***	$\Xi(1820)$	3/2 ⁻ ***	$\Lambda_c(2880)^+$	5/2 ⁺ ***
$N(1650)$	1/2 ⁻ ***	$\Delta(1900)$	1/2 ⁻ **	$\Sigma(1580)$	3/2 ⁻ *	$\Xi(1950)$	***	$\Lambda_c(2940)^+$	***
$N(1675)$	5/2 ⁻ ***	$\Delta(1905)$	5/2 ⁺ ****	$\Sigma(1620)$	1/2 ⁻ *	$\Xi(2030)$	$\geq \frac{5}{2}$ ***	$\Sigma_c(2455)$	1/2 ⁻ ****
$N(1680)$	5/2 ⁺ ***	$\Delta(1910)$	1/2 ⁺ ****	$\Sigma(1620)$	1/2 ⁻ *	$\Xi(2120)$	*	$\Sigma_c(2520)$	3/2 ⁻ ***
$N(1685)$	*	$\Delta(1920)$	3/2 ⁺ ***	$\Sigma(1660)$	1/2 ⁺ ***	$\Xi(2250)$	**	$\Sigma_c(2800)$	***
$N(1700)$	3/2 ⁻ ***	$\Delta(1930)$	5/2 ⁻ ***	$\Sigma(1670)$	3/2 ⁻ ***	$\Xi(2250)$	**	Ξ_c^+	1/2 ⁻ ***
$N(1710)$	1/2 ⁺ ***	$\Delta(1940)$	3/2 ⁻ **	$\Sigma(1690)$	***	$\Xi(2370)$	**	Ξ_c^+	1/2 ⁻ ***
$N(1720)$	3/2 ⁻ ***	$\Delta(1950)$	7/2 ⁺ ****	$\Sigma(1730)$	3/2 ⁺ *	$\Xi(2500)$	*	Ξ_c^0	1/2 ⁻ ***
$N(1860)$	5/2 ⁺ **	$\Delta(2000)$	5/2 ⁺ **	$\Sigma(1750)$	1/2 ⁻ ***	Ω^-	3/2 ⁺ ****	$\Xi_c(2645)$	3/2 ⁻ ***
$N(1875)$	3/2 ⁻ ***	$\Delta(2150)$	1/2 ⁻ *	$\Sigma(1770)$	1/2 ⁺ *	$\Omega(2250)^-$	***	$\Xi_c(2790)$	1/2 ⁻ ***
$N(1880)$	1/2 ⁺ **	$\Delta(2200)$	7/2 ⁻ *	$\Sigma(1775)$	5/2 ⁻ ***	$\Omega(2380)^-$	***	$\Xi_c(2815)$	3/2 ⁻ ***
$N(1895)$	1/2 ⁻ **	$\Delta(2300)$	9/2 ^{+*} **	$\Sigma(1840)$	3/2 ⁺ *	$\Omega(2470)^-$	**	$\Xi_c(2930)$	*
$N(1900)$	3/2 ⁺ ***	$\Delta(2350)$	5/2 ⁻ *	$\Sigma(1880)$	1/2 ⁺ **	$\Xi_c(2980)$	***	$\Xi_c(3055)$	***
$N(1990)$	7/2 ⁺ **	$\Delta(2390)$	7/2 ⁺ *	$\Sigma(1900)$	1/2 ⁻ *	$\Xi_c(2980)$	***	$\Xi_c(3080)$	***
$N(2000)$	5/2 ⁺ **	$\Delta(2400)$	9/2 ⁻ **	$\Sigma(1915)$	5/2 ⁺ ****	$\Xi_c(3123)$	*	$\Xi_c(3123)$	*
$N(2040)$	3/2 ⁺ *	$\Delta(2420)$	11/2 ⁺ ****	$\Sigma(1940)$	3/2 ⁺ *	Ω_c^0	1/2 ⁺ ***	$\Omega_c(2770)^0$	3/2 ⁻ ***
$N(2060)$	5/2 ⁻ **	$\Delta(2750)$	13/2 ⁻ **	$\Sigma(1940)$	3/2 ⁻ ***	Ξ_c^+	*	$\Xi_c(2770)^0$	1/2 ⁺ ***
$N(2100)$	1/2 ⁺ *	$\Delta(2950)$	15/2 ⁺ **	$\Sigma(2000)$	1/2 ⁻ *	Ξ_c^0	*	$K(3100)$? (? ?)
$N(2120)$	3/2 ⁻ **	$\Sigma(2030)$	7/2 ⁻ ****	$\Xi_c(2070)$	5/2 ⁺ *	Ξ_c^0	*	$K(3100)$? (? ?)
$N(2190)$	7/2 ⁻ ***	Λ	1/2 ⁺ ****	$\Sigma(2250)$	***	Ξ_c^0	1/2 ⁻ ***	$\Xi_c(2770)^0$	3/2 ⁻ ***
$N(2220)$	9/2 ⁻ ***	$\Lambda(1405)$	1/2 ⁻ ***	$\Sigma(2080)$	3/2 ⁻ **	Ξ_c^0	*	$K(3100)$? (? ?)
$N(2250)$	9/2 ⁻ ***	$\Lambda(1520)$	3/2 ⁻ ***	$\Sigma(2100)$	7/2 ⁻ *	Λ_b^0	1/2 ⁺ ***	$\Xi_c(2770)^0$	1/2 ⁻ ***
$N(2300)$	1/2 ⁺ **	$\Lambda(1600)$	1/2 ⁺ ***	$\Sigma(2250)$	***	$\Lambda_b(5912)^0$	1/2 ⁻ ***	$\Xi_c(2770)^0$	3/2 ⁻ ***
$N(2570)$	5/2 ⁻ **	$\Lambda(1670)$	1/2 ⁻ ***	$\Sigma(2455)$	**	$\Lambda_b(5920)^0$	3/2 ⁻ ***	$\Xi_c(2770)^0$	1/2 ⁻ ***
$N(2600)$	11/2 ⁻ ***	$\Lambda(1690)$	3/2 ⁻ ***	$\Sigma(2620)$	**	Σ_b	1/2 ⁻ ***	$\Xi_c(2770)^0$	3/2 ⁻ ***
$N(2700)$	13/2 ⁻ **	$\Lambda(1710)$	1/2 ⁺ *	$\Sigma(3000)$	*	Σ_b^+	3/2 ⁻ ***	$\Xi_c(2770)^0$	1/2 ⁻ ***
$\Lambda(1800)$	1/2 ⁻ ***	$\Sigma(3170)$	*	Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***
$\Lambda(1810)$	1/2 ⁺ ***			Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***
$\Lambda(1820)$	5/2 ⁺ ***			Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***
$\Lambda(1830)$	5/2 ⁻ ***			Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***
$\Lambda(1890)$	3/2 ⁺ ***			Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***
$\Lambda(2000)$	*			Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***
$\Lambda(2020)$	7/2 ⁺ *			Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***
$\Lambda(2050)$	3/2 ⁻ *			Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***
$\Lambda(2100)$	7/2 ⁻ ***			Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***
$\Lambda(2110)$	5/2 ⁺ ***			Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***
$\Lambda(2325)$	3/2 ⁻ *			Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***
$\Lambda(2350)$	9/2 ⁺ ***			Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***
$\Lambda(2585)$	**			Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***	Ξ_b^0	1/2 ⁻ ***



PDG2015 : <http://pdg.lbl.gov/>

LIGHT UNFLAVORED ($S = C = B = 0$)		STRANGE ($S = \pm 1, C = B = 0$)		CHARMED, STRANGE ($C = S = \pm 1$)		$\Xi_c(F^C)$
$F(F^C)$	$J(F^C)$	$F(F^C)$	$J(F^C)$	$F(F^C)$	$J(F^C)$	$\Xi_c(F^C)$
$\bullet \pi^\pm$	1 ⁻ (0 ⁻)	$\bullet \phi(1680)$	0 ⁻ (1 ⁻)	$\bullet K^\pm$	1/2(0 ⁻)	$\bullet D_s^+$
$\bullet \eta^0$	1 ⁻ (0 ⁻)	$\bullet \rho(1690)$	1 ⁻ (3 ⁻)	$\bullet K^0$	1/2(0 ⁻)	$\bullet D_s^0$
$\bullet f_0(500)$	0 ⁺ (0 ⁻)	$\bullet \rho(1700)$	1 ⁺ (1 ⁻)	$\bullet K_S^0$	1/2(0 ⁻)	$\bullet D_s^0(2317)^0$
$\bullet \pi(770)$	1 ⁺ (1 ⁻)	$\bullet \rho(1710)$	0 ⁺⁽⁺⁾	$\bullet K_0^0(800)$	1/2(0 ⁺)	$\bullet D_s(2460)^0$
$\bullet \omega(782)$	0 ⁻ (1 ⁻)	$\bullet \eta(1760)$	0 ^{+(0⁻)}	$\bullet K'(892)$	1/2(1 ⁻)	$\bullet D_s(2536)^0$
$\bullet \eta(958)$	0 ^{+(0⁻)}	$\bullet \pi(1800)$	1 ⁻ (0 ⁻)	$\bullet K_{1270}(1270)$	1/2(0 ⁺)	$\bullet D_s(2700)^0$
$\bullet f_0(980)$	0 ^{+(0⁻)}	$\bullet \rho(1810)$	0 ⁺⁽⁺⁾	$\bullet K_1(1400)$	1/2(2 ⁻)	$\bullet D_s^*(2860)^0$
$\bullet \chi_c(980)$	1 ^{-(0⁻)}	$\bullet \chi_c(1835)$??(?)	$\bullet K'(1410)$	1/2(2 ⁻)	$\bullet D_s(3040)^0$
$\bullet \chi_c(1020)$	0 ^{-(1⁻)}	$\bullet \chi_c(1840)$??(?)	$\bullet K_1(1430)$	1/2(2 ⁺)	
$\bullet h_1(1170)$	0 ⁻ (1 ⁻)	$\bullet \phi(1850)$	0 ^{-(3⁻)}	$\bullet K'(1460)$	1/2(0 ⁻)	
$\bullet b_1(1235)$	1 ^{+(1⁻)}	$\bullet \pi(1870)$	0 ^{+(2⁻)}	$\bullet K_{1770}(1770)$	1/2(0 ⁻)	
$\bullet \chi_c(1260)$	1 ^{-(1⁻)}	$\bullet \pi(1880)$	1 ^{-(2⁻)}	$\bullet K_2(1780)(1780)$	1/2(3 ⁻)	
$\bullet f_0(1270)$	0 ^{+(2⁻)}	$\bullet \rho(1900)$	1 ^{-(1⁻)}	$\bullet K_2(1820)(1820)$	1/2(2 ⁻)	
$\bullet f_0(1285)$	0 ^{+(0⁻)}	$\bullet \chi_c(1910)$??(?)	$\bullet K_2(1830)(1830)$	1/2(2 ⁺)	
$\bullet \chi_c(1300)$	1 ^{-(0⁻)}	$\bullet \rho(1920)$	0 ^{+(1⁻)}	$\bullet K_2(1860)(1860)$	1/2(2 ⁺)	
$\bullet \chi_c(1320)$	1 ^{-(2⁻)}	$\bullet \chi_c(1930)$	0 ^{+(0⁻)}	$\bullet K_2(1970)(1970)$	1/2(0 ⁻)	
$\bullet f_0(1370)$	0 ^{+(0⁻)}	$\bullet \rho(1940)$	0 ^{+(0⁻)}	$\bullet K_2(2050)(2050)$	1/2(2 ⁻)	
$\bullet h_1(1380)$	0 ^{-(1⁻)}	$\bullet \chi_c(1940)$	1 ^{-(4⁻)}	$\bullet K_2(2250)(2250)$	1/2(2 ⁻)	
$\bullet \pi_1(1400)$	1 ^{-(1⁻)}	$\bullet \chi_c(1950)$	0 ^{+(4⁻)}	$\bullet K_2(2320)(2320)$	1/2(3 ⁻)	
$\bullet \pi_1(1405)$	0 ^{+(0⁻)}	$\bullet \chi_c(1960)$	0 ^{+(7⁻)}	$\bullet K_2(2380)(2380)$	1/2(2 ⁻)	
$\bullet \chi_c(1475)$	0 ^{+(0⁻)}	$\bullet \chi_c(2020)$	0 ^{+(0⁻)}	$\bullet K_2(2500)(2500)$	1/2(4 ⁻)	
$\bullet f_0(1500)$	0 ^{+(0⁻)}	$\bullet \eta(2225)$	0 ^{+(0⁻)}	$\bullet K_2(2710)(2710)$??(?)	
$\bullet f_0(1510)$	0 ^{+(1⁻)}	$\bullet \rho(2250)$	1 ^{-(3⁻)}	$\bullet B_s^0(5830)^0$	0(1 ⁺)	
$\bullet f_0(1525)$	0 ^{+(2⁻)}	$\bullet f_0(2300)$	0 ^{+(4⁻)}	$\bullet B_s^0(5840)^0$	0(2 ⁺)	
$\bullet f_0(1565)$	0 ^{+(2⁻)}	$\bullet f_0(2300)$	0 ^{+(0⁻)}	$\bullet B_s^*(5880)$??(?)	
$\bullet \rho(1570)$	1 ^{+(1⁻)}	$\bullet f_0(2300)$	0 ^{+(0⁻)}	$\bullet D^*(2010)^0$	1/2(2 ⁻)	
$\bullet f_0(2340)$	0 ^{+(2⁻)}	$\bullet \rho(2350)$	1 ^{+(1⁻)}	$\bullet D_1(2400)^0$	1/2(0 ⁺)	
$\bullet \pi_1(1600)$	1 ^{-(1⁻)}	$\bullet \rho(2350)$	1 ^{+(6⁻)}	$\bullet D_2(2400)^{\pm}$	1/2(0 ⁺)	
$\bullet \chi_c(1640)$	1 ^{-(1⁻)}	$\bullet \chi_c(2450)$	1 ^{-(6⁻)}	$\bullet D_3(2420)^0$	1/2(2 ⁺)	
$\bullet f_0(1645)$	0 ^{+(2⁻)}	$\bullet \chi_c(2510)$	0 ^{+(6⁻)}	$\bullet D_4(2420)^{\pm}$	1/2(2 ⁺)	
$\bullet \pi_2(1650)$	0 ^{-(1⁻)}			$\bullet D_5(2420)^{\pm}$	1/2(2 ⁺)	
$\bullet \pi_3(1670)$	0 ^{-(3⁻)}			$\bullet D_6(2430)^0$	1/2(2 ⁺)	
$\bullet \pi_2(1670)$	1 ^{-(2⁻)}			$\bullet D_7(2430)^0$	1/2(2 ⁺)	
$\bullet \Omega_b^0$	1/2 ⁺ ***			$\bullet D_8(2460)^0$	1/2(2 ⁺)	
$\bullet \Xi_b^0$	1/2 ⁻ ***			$\bullet D_9(2460)^{\pm}$	1/2(2 ⁺)	
$\bullet \Xi_b^0$	1/2 ⁻ ***			$\bullet D_10(2460)^0$	1/2(2 ⁻)	
$\bullet \Xi_b^0$	1/2 ⁻ ***			$\bullet D_11(2460)^{\pm}$	1/2(2 ⁺)	
$\bullet \Xi_b^0$	1/2 ⁻ ***			$\bullet D_12(2460)^0$	1/2(2 ⁻)	
$\bullet \Xi_b^0$	1/2 ⁻ ***			$\bullet D_13(2460)^{\pm}$	1/2(2 ⁺)	
$\bullet \Xi_b^0$	1/2 ⁻ ***			$\bullet D_14(2460)^0$	1/2(2 ⁻)	
$\bullet \Xi_b^0$	1/2 ⁻ ***			$\bullet D_15(2460)^{\pm}$	1/2(2 ⁺)	
$\bullet \Xi_b^0$	1/2 ⁻ ***			$\bullet D_16(2460)^0$	1/2(2 ⁻)	
$\bullet \Xi_b^0$	1/2 ⁻ ***			$\bullet D_17(2460)^{\pm}$	1/2(2 ⁺)	
$\bullet \Xi_b^0$	1/2 ⁻ ***			$\bullet D_18(2460)^0$	1/2(2 ⁻)	
$\bullet \Xi_b^0$	1/2 ⁻ ***			$\bullet D_19(2460)^{\pm}$	1/2(2 ⁺)	
$\bullet \Xi_b^0$	1/2 ⁻ ***			$\bullet D_20(2460)^0$	1/2(2 ⁻)	
$\bullet \Xi_b^0$	1/2 ⁻ ***			$\bullet D_21(2460)^{\pm}$	1/2(2 ⁺)	
$\bullet \Xi_b^0$	1/2 ⁻ ***			$\bullet D_22(2460)^0$	1/2(2 ⁻)	
$\bullet \Xi_b^0$	1/2 ^{-</sup}					

Difficulty of resonances

Resonance as an “eigenstate” of Hamiltonian

- complex energy

G. Gamow, Z. Phys. 51, 204 (1928)

Zur Quantentheorie des Atomkernes.

Von G. Gamow, z. Zt. in Göttingen.

Mit 5 Abbildungen. (Eingegangen am 2. August 1928.)

Um diese Schwierigkeit zu überwinden, müssen wir annehmen, daß die Schwingungen gedämpft sind, und E komplex setzen:

$$E = E_0 + i \frac{\hbar \lambda}{4 \pi},$$

wo E_0 die gewöhnliche Energie ist und λ das Dämpfungsdekkrement (Zerfallskonstante). Dann sehen wir aber aus den Relationen (2a) und (2b),

- Wave function is not normalized ($\text{Im } k < 0$).

$$\langle R | R \rangle = \int d\mathbf{r} |\psi_R(\mathbf{r})|^2 \sim \int_0^\infty dr e^{-2\text{Im}[k]r} \rightarrow \infty$$

Bi-orthogonal basis (Gamow vectors): normalizable!

N. Hokkyo, Prog. Theor. Phys. 33, 1116 (1965)

T. Berggren, Nucl. Phys. A 109, 265 (1968)


$$| \tilde{R} \rangle = | R^* \rangle, \quad | \langle \tilde{R} | R \rangle | = \left| \int d\mathbf{r} [\psi_R(\mathbf{r})]^2 \right| < \infty$$

- Complex expectation value (e.g. $\langle r^2 \rangle$) —> interpretation?

Resonances in quantum mechanics



1) Potential (shape) resonance

- single channel (P)
- potential barrier : $E > 0$
- unstable via tunneling
- (composite of P -channel)



2) Feshbach resonance

- coupled channel ($P+Q$)
- bound state of Q : $E_Q < 0$, $E_P > 0$
- unstable via transition
- (“elementary”: other than P)




Compositeness of hadrons

- Internal structure of excited hadrons
- Weak binding relation for stable bound states

S. Weinberg, Phys. Rev. 137, B672 (1965)

Compositeness X
threshold channel



or

“Elementariness” Z
other contributions



observables

- Study compositeness of hadron resonances by generalizing weak binding relation to unstable states with effective field theory.

Weak binding relation for stable states

Compositeness of s-wave weakly bound state ($R \gg R_{\text{typ}}$)

S. Weinberg, Phys. Rev. 137, B672 (1965);

T. Hyodo, Int. J. Mod. Phys. A 28, 1330045 (2013)

$$a_0 = R \left\{ \frac{2X}{1+X} + \mathcal{O}\left(\frac{R_{\text{typ}}}{R}\right) \right\}, \quad r_e = R \left\{ \frac{X-1}{X} + \mathcal{O}\left(\frac{R_{\text{typ}}}{R}\right) \right\}$$

a_0 : scattering length, r_e : effective range

$R = (2\mu B)^{-1/2}$: radius of wave function

R_{typ} : length scale of interaction

X: probability of finding composite component

- deuteron is NN composite ($a_0 \sim R \gg r_e$) $\rightarrow X \sim 1$
- internal structure from **observable**
- no nuclear force potential / wavefunction of deuteron

Problem: applicable only for stable states

Effective field theory

Low-energy scattering with near-threshold bound state


- Nonrelativistic EFT with contact interaction

D.B. Kaplan, Nucl. Phys. B494, 471 (1997)

Y. Kamiya, T. Hyodo, Phys. Rev. C93, 035203 (2016)

$$H_{\text{free}} = \int d\mathbf{r} \left[\frac{1}{2M} \nabla \psi^\dagger \cdot \nabla \psi + \frac{1}{2m} \nabla \phi^\dagger \cdot \nabla \phi + \frac{1}{2M_0} \nabla B_0^\dagger \cdot \nabla B_0 + \nu_0 B_0^\dagger B_0 \right],$$

$$H_{\text{int}} = \int d\mathbf{r} \left[g_0 \left(B_0^\dagger \phi \psi + \psi^\dagger \phi^\dagger B_0 \right) + v_0 \psi^\dagger \phi^\dagger \phi \psi \right]$$



- **cutoff** : $\Lambda \sim 1/R_{\text{typ}}$ (**interaction range of microscopic theory**)
- At low momentum $p \ll \Lambda$, interaction \sim **contact**

Compositeness and “elementariness”

Eigenstates

$$H_{\text{free}} |B_0\rangle = \nu_0 |B_0\rangle, \quad H_{\text{free}} |\mathbf{p}\rangle = \frac{\mathbf{p}^2}{2\mu} |\mathbf{p}\rangle \quad \text{free (discrete + continuum)}$$

$$(H_{\text{free}} + H_{\text{int}}) |B\rangle = -B |B\rangle \quad \text{full (bound state)}$$


- normalization of $|B\rangle$ + completeness relation

$$\langle B | B \rangle = 1, \quad 1 = |B_0\rangle\langle B_0| + \int \frac{d\mathbf{p}}{(2\pi)^3} |\mathbf{p}\rangle\langle\mathbf{p}|$$

- projections onto bare states

$$1 = Z + X, \quad Z \equiv |\langle B_0 | B \rangle|^2, \quad X \equiv \int \frac{d\mathbf{p}}{(2\pi)^3} |\langle \mathbf{p} | B \rangle|^2$$

“elementariness” compositeness




Z, X : real and nonnegative \rightarrow interpreted as probability

Weak binding relation

$\Psi\Phi$ scattering amplitude (exact result)

$$f(E) = -\frac{\mu}{2\pi} \frac{1}{[v(E)]^{-1} - G(E)}$$



$$v(E) = v_0 + \frac{g_0^2}{E - \nu_0}, \quad G(E) = \frac{1}{2\pi^2} \int_0^\Lambda dp \frac{p^2}{E - p^2/(2\mu) + i0^+}$$

Compositeness $\times \leftarrow v(E), G(E)$

T. Sekihara, T. Hyodo, D. Jido, PTEP2015, 063D04 (2015)

$$X = \{1 + G^2(-B)v'(-B) [G'(-B)]^{-1}\}^{-1}$$

$1/R=(2\mu B)^{1/2}$ expansion: leading term $\leftarrow X$

$$a_0 = -f(E=0) = R \left\{ \frac{2X}{1+X} + \overline{\mathcal{O}\left(\frac{R_{\text{typ}}}{R}\right)} \right\}$$

renormalization dependent

renormalization independent

If $R \gg R_{\text{typ}}$, correction terms neglected: $X \leftarrow (B, a_0)$

Introduction of decay channel

Introduce decay channel


$$H'_{\text{free}} = \int d\mathbf{r} \left[\frac{1}{2M'} \nabla \psi'^{\dagger} \cdot \nabla \psi' - \nu_{\psi} \psi'^{\dagger} \psi' + \frac{1}{2m'} \nabla \phi'^{\dagger} \cdot \nabla \phi' - \nu_{\phi} \phi'^{\dagger} \phi' \right],$$

$$H'_{\text{int}} = \int d\mathbf{r} \left[g'_0 \left(B_0^{\dagger} \phi' \psi' + \psi'^{\dagger} \phi'^{\dagger} B_0 \right) + v'_0 \psi'^{\dagger} \phi'^{\dagger} \phi' \psi' + v_0^t (\psi^{\dagger} \phi^{\dagger} \phi' \psi' + \psi'^{\dagger} \phi'^{\dagger} \phi \psi) \right],$$

Quasi-bound state: complex eigenvalue

$$H = H_{\text{free}} + H'_{\text{free}} + H_{\text{int}} + H'_{\text{int}}$$

$$H|QB\rangle = E_{QB}|QB\rangle, \quad E_{QB} \in \mathbb{C}$$



Generalized relation: correction term \leftarrow threshold difference

$$a_0 = R \left\{ \frac{2X}{1+X} + \mathcal{O} \left(\left| \frac{R_{\text{typ}}}{R} \right| \right) + \underline{\mathcal{O} \left(\left| \frac{l}{R} \right|^3 \right)} \right\}, \quad R = \frac{1}{\sqrt{-2\mu E_{QB}}} \in \mathbb{C}, \quad l \equiv \frac{1}{\sqrt{2\mu\nu}}$$

[Y. Kamiya, T. Hyodo, Phys. Rev. C93, 035203 \(2016\)](#)

c.f. V. Baru, *et al.*, Phys. Lett. B586, 53 (2004), ...

If $|R| \gg (R_{\text{typ}}, |l|)$ correction terms neglected: $X \leftarrow (E_{QB}, a_0)$

Recent developments

Complex X: interpretation?

$$\tilde{Z} = \frac{1 - |X| + |Z|}{2}, \quad \tilde{X} = \frac{1 - |Z| + |X|}{2} \quad \tilde{Z} + \tilde{X} = 1, \quad \tilde{Z}, \tilde{X} \in [0, 1]$$

compositeness when $U = |Z| + |X| - 1$ is small

[Y. Kamiya, T. Hyodo, Phys. Rev. C93, 035203 \(2016\)](#)

CDD pole contribution (Pade approximant for inverse amp.)

$$X_{\text{Pade}} = \left[1 - \frac{4R(a_0 - R)^2}{a_0^2 r_e} \right]^{-1}$$

[Y. Kamiya, T. Hyodo, arXiv:1607.01899\[hep-ph\]](#)

Talk by Y. Kamiya

Formulation with finite volume effect

- difficulty of resonance \leftrightarrow continuum (c.f. plane wave)
- Finite volume: only discrete eigenstates

$$X_{FV}(L) \in [0, 1], \quad \lim_{L \rightarrow \infty} X_{FV}(L)?$$

[Y. Tsuchida, T. Hyodo, in preparation](#)

Talk by Y. Tsuchida

Application

Generalized weak binding relation $X \leftarrow (E_{QB}, a_0)$

$$a_0 = R \left\{ \frac{2X}{1+X} + \mathcal{O}\left(\left|\frac{R_{typ}}{R}\right|\right) + \mathcal{O}\left(\left|\frac{l}{R}\right|^3\right) \right\}, \quad R = \frac{1}{\sqrt{-2\mu E_{QB}}}, \quad l \equiv \frac{1}{\sqrt{2\mu\nu}}$$

- $\Lambda(1405)$ (higher) pole position and $\bar{K}N$ scattering length

Y. Ikeda, T. Hyodo, W. Weise, PLB 706, 63 (2011); NPA 881 98 (2012), ...

- $E_{QB} = -10 - 26i$ MeV $\rightarrow |R| \sim 2$ fm \rightarrow small correction term

$\left|\frac{R_{typ}}{R}\right| \lesssim 0.12, \quad \left|\frac{l}{R}\right|^3 \lesssim 0.16$ energy difference from $\pi\Sigma$
↑
↓
vector meson exchange

Ref.	E_{QB} (MeV)	a_0 (fm)	$X_{\bar{K}N}$	$\tilde{X}_{\bar{K}N}$	U	$ r_e/a_0 $
[43]	$-10 - i26$	$1.39 - i0.85$	$1.2 + i0.1$	1.0	0.5	0.2
[44]	$-4 - i8$	$1.81 - i0.92$	$0.6 + i0.1$	0.6	0.0	0.7
[45]	$-13 - i20$	$1.30 - i0.85$	$0.9 - i0.2$	0.9	0.1	0.2
[46]	$2 - i10$	$1.21 - i1.47$	$0.6 + i0.0$	0.6	0.0	0.7
[46]	$-3 - i12$	$1.52 - i1.85$	$1.0 + i0.5$	0.8	0.6	0.4

↑
systematic error
↓

$\Lambda(1405)$ is $\bar{K}N$ composite \leftarrow observables

Summary



Compositeness of near-threshold bound state can be determined only by observables.

S. Weinberg, Phys. Rev. 137, B672 (1965)




Weak binding relation can be generalized to unstable states with effective field theory.

$$a_0 = R \left\{ \frac{2X}{1+X} + \mathcal{O}\left(\left|\frac{R_{typ}}{R}\right|\right) + \mathcal{O}\left(\left|\frac{l}{R}\right|^3\right) \right\}, \quad R = \frac{1}{\sqrt{-2\mu E_{QB}}}, \quad l \equiv \frac{1}{\sqrt{2\mu\nu}}$$



Precise determination of the pole position and scattering length shows that $\Lambda(1405)$ is dominated by $\bar{K}N$ composite component.



[Y. Kamiya, T. Hyodo, Phys. Rev. C93, 035203 \(2016\), arXiv:1607.01899\[hep-ph\]](#)
[Y. Tsuchida, T. Hyodo, in preparation](#)