

閾値近傍状態の構造について



Tetsuo Hyodo

Tokyo Metropolitan Univ.


2020, Mar. 17th 1

Classification of hadrons

PDG2018 : <http://pdg.lbl.gov/>


Usual hadrons ← quarks

p	$1/2^+$	****	$\Delta(1232)$	$3/2^+$	****	Σ^+	$1/2^+$	****	Ξ^0	$1/2^+$	****	Λ_c^+	$1/2^+$	****
n	$1/2^+$	****	$\Delta(1600)$	$3/2^+$	***	Σ^0	$1/2^+$	****	Ξ^-	$1/2^+$	****	$\Lambda_c(2595)^+$	$1/2^-$	***
$N(1440)$	$1/2^+$	****	$\Delta(1620)$	$1/2^-$	****	Σ^-	$1/2^+$	****	$\Xi(1530)$	$3/2^+$	****	$\Lambda_c(2625)^+$	$3/2^-$	***
$N(1520)$	$3/2^-$	****	$\Delta(1700)$	$3/2^-$	****	$\Sigma(1385)$	$3/2^+$	****	$\Xi(1620)$	*	****	$\Lambda_c(2765)^+$	*	****
$N(1535)$	$1/2^-$	****	$\Delta(1750)$	$1/2^+$	*	$\Sigma(1480)$	*	****	$\Xi(1690)$	***	****	$\Lambda_c(2880)^+$	$5/2^+$	***
$N(1650)$	$1/2^-$	****	$\Delta(1900)$	$1/2^-$	**	$\Sigma(1560)$	**	****	$\Xi(1820)$	$3/2^-$	***	$\Lambda_c(2940)^+$	***	****
$N(1675)$	$5/2^-$	****	$\Delta(1905)$	$5/2^+$	****	$\Sigma(1580)$	$3/2^-$	**	$\Xi(1950)$	$\geq 5/2^+$	****	$\Sigma_c(2455)$	$1/2^+$	****
$N(1680)$	$5/2^+$	****	$\Delta(1910)$	$1/2^+$	****	$\Sigma(1620)$	$1/2^-$	*	$\Xi(2030)$	$\geq 5/2^+$	****	$\Sigma_c(2520)$	$3/2^+$	****
$N(1685)$	*	****	$\Delta(1920)$	$3/2^+$	***	$\Sigma(1660)$	$1/2^+$	***	$\Xi(2120)$	**	****	$\Sigma_c(2800)$	***	****
$N(1700)$	$3/2^-$	***	$\Delta(1930)$	$5/2^-$	***	$\Sigma(1670)$	$3/2^-$	****	$\Xi(2250)$	**	****	Ξ_c^+	$1/2^+$	****
$N(1710)$	$1/2^+$	***	$\Delta(1940)$	$3/2^-$	**	$\Sigma(1690)$	**	****	$\Xi(2370)$	**	****	Ξ_c^0	$1/2^+$	****
$N(1720)$	$3/2^+$	****	$\Delta(1950)$	$7/2^+$	****	$\Sigma(1730)$	$3/2^+$	**	$\Xi(2500)$	*	****	Ξ_c^-	$1/2^+$	****
$N(1860)$	$5/2^+$	**	$\Delta(2000)$	$5/2^+$	**	$\Sigma(1750)$	$1/2^-$	***	Ω^-	$3/2^+$	****	$\Xi_c(2645)$	$3/2^+$	****
$N(1875)$	$3/2^-$	***	$\Delta(2150)$	$1/2^-$	*	$\Sigma(1770)$	$1/2^+$	**	$\Omega(2250)$	****	****	$\Xi_c(2790)$	$1/2^-$	****
$N(1880)$	$1/2^+$	**	$\Delta(2200)$	$7/2^-$	*	$\Sigma(1775)$	$5/2^-$	****	$\Omega(2380)$	**	****	$\Xi_c(2815)$	$3/2^-$	****
$N(1895)$	$1/2^-$	**	$\Delta(2300)$	$9/2^+$	**	$\Sigma(1840)$	$3/2^+$	**	$\Omega(2470)$	**	****	$\Xi_c(2930)$	*	****
$N(1900)$	$3/2^+$	***	$\Delta(2350)$	$5/2^-$	**	$\Sigma(1880)$	$1/2^+$	**				$\Xi_c(2980)$	***	****
$N(1990)$	$7/2^+$	**	$\Delta(2390)$	$7/2^+$	*	$\Sigma(1900)$	$1/2^-$	*				$\Xi_c(3055)$	***	****
$N(2000)$	$5/2^+$	**	$\Delta(2400)$	$9/2^-$	**	$\Sigma(1915)$	$5/2^+$	****				$\Xi_c(3080)$	***	****
$N(2040)$	$3/2^+$	*	$\Delta(2420)$	$11/2^+$	****	$\Sigma(1940)$	$3/2^+$	**				$\Xi_c(3123)$	*	****
$N(2060)$	$5/2^-$	**	$\Delta(2750)$	$13/2^-$	**	$\Sigma(1940)$	$3/2^-$	***				Ω_c^0	$1/2^+$	****
$N(2100)$	$1/2^+$	*	$\Delta(2950)$	$15/2^+$	**	$\Sigma(2000)$	$1/2^-$	*				$\Omega_c(2770)^0$	$3/2^+$	****
$N(2120)$	$3/2^-$	**				$\Sigma(2030)$	$7/2^+$	****				Ξ_{cc}^+	*	****
$N(2190)$	$7/2^-$	****	$\Lambda(1405)$	$1/2^-$	****	$\Sigma(2070)$	$5/2^+$	*						
$N(2220)$	$9/2^+$	****	$\Lambda(1520)$	$3/2^-$	****	$\Sigma(2080)$	$3/2^+$	**						
$N(2250)$	$9/2^-$	****	$\Lambda(1600)$	$1/2^+$	****	$\Sigma(2100)$	$7/2^-$	*						
$N(2300)$	$1/2^+$	**	$\Lambda(1670)$	$1/2^-$	****	$\Sigma(2250)$	***	****				Λ_b^0	$1/2^+$	****
$N(2570)$	$5/2^-$	**	$\Lambda(1690)$	$3/2^-$	****	$\Sigma(2455)$	**	****				$\Lambda_b(5912)^0$	$1/2^-$	****
$N(2600)$	$11/2^-$	***	$\Lambda(1710)$	$1/2^+$	*	$\Sigma(2620)$	**	****				$\Lambda_b(5920)^0$	$3/2^-$	****
$N(2700)$	$13/2^+$	**	$\Lambda(1800)$	$1/2^-$	***	$\Sigma(3000)$	*	****				$\Lambda_b(5955)^0$	$3/2^+$	****
			$\Lambda(1810)$	$1/2^+$	***	$\Sigma(3170)$	*	****				Ω_b^-	$1/2^+$	****
			$\Lambda(1820)$	$5/2^+$	****									
			$\Lambda(1830)$	$5/2^-$	****									
			$\Lambda(1890)$	$3/2^+$	****									
			$\Lambda(2000)$	*	****									
			$\Lambda(2020)$	$7/2^+$	*									
			$\Lambda(2050)$	$3/2^-$	*									
			$\Lambda(2100)$	$7/2^-$	****									
			$\Lambda(2110)$	$5/2^+$	***									
			$\Lambda(2325)$	$3/2^-$	*									
			$\Lambda(2350)$	$9/2^+$	**									
			$\Lambda(2585)$	**	****									



~ 150 baryons

LIGHT UNFLAVORED (S = C = B = 0)		STRANGE (S = ±1, C = B = 0)		CHARMED, STRANGE (C = S = ±1)		CC F _c (F _c)	
F(F _c)		F(F _c)		F(F _c)		F(F _c)	
π^\pm	$1^-(0^-)$	$\rho(1680)$	$0^-(1^-)$	K^\pm	$1/2(0^-)$	D_s^\pm	$0(0^-)$
π^0	$1^-(0^-)$	$\rho(1690)$	$1^+(3^-)$	K^0	$1/2(0^-)$	D_s^\pm	$0(?)^2$
η	$0^+(0^+)$	$\rho(1700)$	$1^+(1^-)$	K_S^0	$1/2(0^-)$	$D_{s1}(2317)$	$0(0^-)$
$\eta(500)$	$0^+(0^+)$	$a_2(1700)$	$1^-(2^+)$	K_L^0	$1/2(0^-)$	$D_{s1}(2460)^+$	$0(1^+)$
$\rho(770)$	$1^+(1^-)$	$\omega(1710)$	$0^+(0^+)$	$K_S^*(800)$	$1/2(0^+)$	$D_{s1}(2536)^+$	$0(1^+)$
$\omega(782)$	$0^+(0^+)$	$\eta(1760)$	$0^+(0^+)$	$K^*(892)$	$1/2(1^-)$	$D_{s2}(2573)$	$0(?)^2$
$\eta(958)$	$0^+(0^+)$	$\pi(1800)$	$1^-(0^+)$	$K_1(1270)$	$1/2(1^+)$	$D_{s1}(2700)^+$	$0(1^-)$
$\phi(930)$	$0^+(0^+)$	$f_2(1810)$	$0^+(2^+)$	$K_1(1400)$	$1/2(1^+)$	$D_{s1}(2860)^+$	$0(?)^2$
$a_0(980)$	$1^-(0^+)$	$X(1835)$	$?^2(?)^+$	$K^*(1410)$	$1/2(1^+)$	$D_{s1}(3040)^+$	$0(?)^2$
$\phi(1020)$	$0^-(1^-)$	$X(1840)$	$?^2(?)^+$	$K_S^*(1430)$	$1/2(0^+)$		
$h(1170)$	$0^-(1^+)$	$\omega_3(1850)$	$0^-(3^-)$	$K_S^*(1430)$	$1/2(0^+)$	BOTTOM (B = ±1)	
$b_1(1235)$	$1^+(1^+)$	$\eta_2(1875)$	$0^+(2^-)$	$K(1460)$	$1/2(0^+)$	B^\pm	$1/2(0^-)$
$a_1(1260)$	$1^-(1^+)$	$\omega_3(1880)$	$1^-(2^+)$	$K_2(1580)$	$1/2(2^-)$	B^0	$1/2(0^-)$
$f_2(1270)$	$0^+(2^+)$	$\pi_2(1900)$	$1^-(1^-)$	$K(1630)$	$1/2(?)^+$	B^+ / B^0	ADMIXTURE
$f_1(1285)$	$0^+(1^+)$	$f_1(1910)$	$0^+(2^+)$	$K_1(1650)$	$1/2(1^+)$	$B^+ / B^0 / B_S^0 / B_{sbaryon}$	ADMIXTURE
$\eta(1295)$	$0^+(0^+)$	$f_1(1950)$	$0^+(2^+)$	$K^*(1680)$	$1/2(1^-)$	V_{cb} and V_{cb}	CKM Matrix Elements
$\pi(1300)$	$1^-(0^+)$	$f_3(1990)$	$1^+(3^-)$	$K^*(1770)$	$1/2(2^-)$	B^*	$1/2(1^-)$
$\phi_2(1320)$	$1^-(2^+)$	$f_2(2010)$	$0^+(2^+)$	$K_S^*(1780)$	$1/2(3^-)$	$B_1(5721)^+$	$1/2(1^+)$
$\phi_3(1370)$	$0^+(0^+)$	$f_0(2020)$	$0^+(0^+)$	$K_S^*(1820)$	$1/2(2^-)$	$B_2(5732)$	$?^2(?)^+$
$h(1380)$	$?^-(1^+)$	$a_0(2040)$	$1^-(4^+)$	$K(1830)$	$1/2(0^+)$	$B_3(5747)^0$	$1/2(2^+)$
$\omega_3(1400)$	$1^-(1^+)$	$f_0(2050)$	$0^+(4^+)$	$K_1(1850)$	$1/2(0^+)$	$B_3(5747)^0$	$1/2(2^+)$
$\eta(1405)$	$0^+(0^+)$	$\pi_2(2100)$	$1^-(2^+)$	$K_S^*(1980)$	$1/2(2^+)$	$B_3(5747)^0$	$1/2(2^+)$
$f_1(1420)$	$0^+(1^+)$	$f_0(2120)$	$0^+(0^+)$	$K_4^*(2045)$	$1/2(4^+)$	$B_3(5747)^0$	$1/2(2^+)$
$\omega(1420)$	$0^-(1^-)$	$f_2(2150)$	$0^+(2^+)$	$K_2(2250)$	$1/2(2^-)$	$B(5970)^+$	$?^2(?)^+$
$f_2(1430)$	$0^+(2^+)$	$\rho(2150)$	$1^+(1^-)$	$K_3(2330)$	$1/2(3^+)$	$B(5970)^0$	$?^2(?)^+$
$a_0(1450)$	$1^-(0^+)$	$\omega(2170)$	$0^-(1^-)$	$K_S^*(2380)$	$1/2(5^-)$	$B(5970)^0$	$?^2(?)^+$
$\phi(1450)$	$1^-(1^-)$	$f_0(2200)$	$0^+(0^+)$	$K_4(2500)$	$1/2(4^-)$		
$\eta(1475)$	$0^+(0^+)$	$f_2(2220)$	$0^+(2^+)$	$K(3100)$	$?^2(?)^+$	BOTTOM, STRANGE (B = ±1, S = ±1)	
$f_0(1500)$	$0^+(0^+)$	$\eta(2225)$	$0^+(0^+)$			B_c^\pm	$0(0^-)$
$f_1(1510)$	$0^+(1^+)$	$\rho_3(2250)$	$1^+(3^-)$	CHARMED (C = ±1)		B_c^\pm	$0(1^-)$
$f_2(1525)$	$0^+(2^+)$	$f_2(2300)$	$0^+(2^+)$	D^\pm	$1/2(0^-)$	$B_{c1}(5830)^0$	$0(1^+)$
$f_3(1565)$	$0^+(2^+)$	$f_0(2330)$	$0^+(0^+)$	D^0	$1/2(0^-)$	$B_{c2}(5840)^0$	$0(2^+)$
$\rho(1570)$	$1^+(1^-)$	$f_2(2340)$	$0^+(0^+)$	$D^*(2007)^0$	$1/2(1^-)$	$B_{c1}(5850)$	$?^2(?)^+$
$h(1595)$	$0^-(1^+)$	$f_3(2340)$	$0^+(2^+)$	$D^*(2010)^+$	$1/2(1^-)$	BOTTOM, CHARMED (B = C = ±1)	
$\pi_1(1600)$	$1^-(1^+)$	$\rho_3(2350)$	$1^+(5^-)$	$D^*(2000)^0$	$1/2(0^+)$	B_c^\pm	$0(0^-)$
$a_1(1640)$	$1^-(1^+)$	$a_0(2450)$	$1^-(6^+)$	$D_1(2400)^+$	$1/2(1^+)$	$B_c(25)^+$	$?^2(?)^+$
$f_2(1640)$	$0^+(2^+)$	$f_0(2510)$	$0^+(6^+)$	$D_1(2430)^0$	$1/2(1^+)$		
$\rho(1645)$	$0^+(2^+)$			$D_1(2460)^+$	$1/2(1^+)$		
$\omega(1650)$	$0^-(1^-)$			$D_2(2460)^0$	$1/2(2^+)$		
$\omega_3(1670)$	$0^-(3^-)$			$D_2(2550)^0$	$1/2(0^-)$		
$\pi_2(1670)$	$1^-(2^+)$			$D_2(2600)^+$	$1/2(2^+)$		
				$D(2600)$	$1/2(?)^+$		
				$D^*(2640)^+$	$1/2(?)^+$		
				$D(2750)$	$1/2(?)^+$		




~ 210 mesons

Can one construct a hadron from hadrons?

Hadron clusters

Hadrons near an **s-wave** two-body threshold



“hadronic molecules” (various flavors, baryon numbers, ...)

Two-body universal physics

Near-threshold s-wave state: **universal physics**

E. Braaten, H.-W. Hammer, Phys. Rept. 428, 259 (2006);

P. Naidon, S. Endo, Rept. Prog. Phys. 80, 056001 (2017)

- scattering length $|a| \gg$ interaction range R_0
- size of (quasi-)bound state $\sim |a|$: loosely bound
- relation with eigenenergy E


$$a(E) \sim \frac{i}{k} = \frac{i}{\sqrt{2\mu E}}$$

vdW

Examples: d , $\Lambda(1405)$, ${}^4\text{He}$ **dimer**

	NN [fm]	$\bar{K}N$ [fm]	${}^4\text{He}$ [a_0]
$a(E)$	4.3	1.2-0.8i	178
a_{emp}	5.1	1.4-0.9i	189
R_0	1.4	0.4	10

strong



${}^4\text{He}$

Wave function

Why is a weakly bound s-wave state spatially large?

- Radial Schrödinger equation


$$-\frac{\hbar^2}{2\mu} \frac{d^2}{dr^2} u_0(r) + V(r)u_0(r) = Eu_0(r), \quad \psi_0(r) = Y_0^0 \frac{u_0(r)}{r}$$

- at large distance ($V(r) = 0$) with zero energy ($E = 0$)

$$u_0(r) = C(r - a), \quad (r > R_0)$$

- scattering length a : intercept of $u_0(r)$

- bound state with $B = 0 \Rightarrow |a| = \infty$



Wave function is **not normalizable**

$$\int d^3r |\psi_0(r)|^2 = \int_0^\infty dr |u_0(r)|^2 = \infty$$

$B = 0$ state is not a bound state (zero-energy resonance)

Consequences

Mean squared radius

$$\langle r^2 \rangle = \int d^3r r^2 |\psi_0(r)|^2 = \int_0^\infty dr r^2 |u_0(r)|^2 = \infty$$

—> size of $B = 0$ state is **infinitely large**

Compositeness X (weight of the composite component)

T. Hyodo, Phys. Rev. C 90, 055208 (2014)

$$\Psi = \begin{pmatrix} \tilde{\psi}_0(p) \\ c \\ \vdots \end{pmatrix} \quad X = \int \frac{d^3p}{(2\pi)^3} |\tilde{\psi}_0(p)|^2 = \int d^3r |\psi_0(r)|^2 \quad \text{infinite}$$

$$Z = |c|^2 + \dots \quad \text{finite}$$

$$X + Z = 1$$

—> $B = 0$ state is **completely composite** ($X = 1$, $Z = 0$)

Weakly bound state ($B \neq 0$, except for fine tuning)

- large spatial size and composite dominance

p-wave state

What about p-wave states?

- Radial Schrödinger equation

$$-\frac{\hbar^2}{2\mu} \frac{d^2}{dr^2} u_1(r) + V(r)u_1(r) + \frac{\hbar^2}{\mu r^2} u_1(r) = E u_1(r), \quad \psi_1(\mathbf{r}) = Y_1^m(\Omega) \frac{u_1(r)}{r}$$

- at large distance ($V(r) = 0$) with zero energy ($E = 0$)

$$u_1(r) = Cr^s, \quad s = 2, -1$$


- bound state with $B = 0$

$$u_1(r) = \frac{C}{r}, \quad (r > R_0)$$

Wave function is **normalizable**

$$\int d^3r |\psi_1(\mathbf{r})|^2 = \int_0^\infty dr |u_1(r)|^2 < \infty$$

$B = 0$ state is a bound state



Consequences (p-wave)

Mean squared radius

$$\langle r^2 \rangle = \int_0^\infty dr r^2 |u_1(r)|^2 = A_{\text{in}} + |C|^2 \int_{R_0}^\infty dr 1 = \infty \quad \leftarrow u_1(r) = \frac{C}{r}$$

→ size of $B = 0$ state is **infinitely large**

Compositeness X

T. Hyodo, Phys. Rev. C 90, 055208 (2014)

$$\Psi = \begin{pmatrix} \tilde{\psi}_1(\mathbf{p}) \\ c \\ \vdots \end{pmatrix} \quad X = \int \frac{d^3p}{(2\pi)^3} |\tilde{\psi}_1(\mathbf{p})|^2 = \int d^3r |\psi_1(\mathbf{r})|^2 \quad \text{finite}$$

$$Z = |c|^2 + \dots \quad \text{finite}$$

$$X + Z = 1$$

→ $B = 0$ can have **any structure** ($0 < X < 1$)

Weakly bound p-wave state ($B \neq 0$, except for fine tuning)

- **large spatial size?**

Higher partial waves

Wave function of a $B = 0$ state with angular momentum ℓ

$$u_\ell(r) = \frac{C}{r^\ell} \quad (r > R_0)$$

- Mean squared radius

$$\langle r^2 \rangle = \int_0^\infty dr r^2 |u_\ell(r)|^2 = A_{\text{in}} + |C|^2 \int_{R_0}^\infty dr r^{2-2\ell} = \infty \quad \text{if } \ell \leq 1$$

—> diverges only for s- and p-waves

Generalized radius

$$\sqrt[n]{\langle r^n \rangle} = \sqrt[n]{\int d^3r r^n |\psi(r)|^2} = \infty \quad \text{if } n - 2\ell \geq 0$$

- s-wave: radius diverges for **all** $n \geq 0$ ← universality


- p-wave: radius diverges for $n \geq 2$

- ℓ -th wave: radius diverges for $n \geq 2\ell$

example: square well potential

Square well potential

$$V(r) = \begin{cases} -V_0 & 0 \leq r \leq b \\ 0 & b < r \end{cases}$$



- s-wave

$$\langle r^2 \rangle = \left[\frac{b}{2} - \frac{\sin(2kb)}{4k} + \frac{\sin^2(kb)}{2\kappa} \right]^{-1} \left[\frac{1}{k^3} \left(\frac{(kb)^3}{6} + \frac{1 - 2(kb)^2}{8} \sin(2kb) - \frac{kb}{4} \cos(2kb) \right) + \frac{\sin^2(kb)}{2\kappa^3} \left[(\kappa b)^2 + \kappa b + \frac{1}{2} \right] \right]$$

$$k = \frac{\sqrt{2\mu(V_0 - B)}}{\hbar}, \quad \kappa = \frac{\sqrt{2\mu B}}{\hbar}$$



- p-wave

$$\langle r^2 \rangle = \left[\frac{b}{2} + \frac{\sin(2kb)}{4k} - \frac{\sin^2(kb)}{k^2 b} + \frac{\kappa (\sin(kb) - kb \cos(kb))^2}{2k^2 (1 + \kappa b)^2} \left(1 + \frac{2}{\kappa b} \right) \right]^{-1}$$

$$\times \left[\frac{1}{k^3} \left(\frac{3kb + (kb)^3}{6} + \frac{-5 + 2(kb)^2}{8} \sin(2kb) + \frac{3kb}{4} \cos(2kb) \right) + \frac{(\sin(kb) - kb \cos(kb))^2}{2\kappa k^2 (1 + \kappa b)^2} \left[(\kappa b)^2 + 3\kappa b + \frac{5}{2} \right] \right]$$

Demonstration

Root mean squared radius $\sqrt{\langle r^2 \rangle}$ v.s. $\kappa = \sqrt{2\mu B/\hbar}$



Both radii diverge in the weak binding limit ($\kappa \rightarrow 0$).

Dependence on b (well width) seems different.

Weak-binding behavior

Behavior near the weak-binding limit

- s-wave

$$\langle r^2 \rangle = \frac{1}{2\kappa^2} + \mathcal{O}(\kappa^{-1})$$



independent of b (well width)

← universality

- p-wave

$$\langle r^2 \rangle = \frac{5b}{6} \frac{1}{\kappa} + \mathcal{O}(\kappa^0)$$

depends on b (well width)
non-universal divergence



Summary



Size and structure of $B = 0$ states

	$\langle r^2 \rangle$	compositeness	$\int d^3r \psi ^2$
s-wave	∞ (universal)	$X = 1$	∞
p-wave	∞ (non-universal)	$0 < X < 1$	finite



Implication: **large** mean squared radius of near-threshold ($B \neq 0$) **p-wave bound states**



What about resonances?

T. Hyodo, in preparation